

Preparing for



Play.

Eastlands Primary School

Juccess in a changing world.

PE Health & Well-being

As we move into winter bugs season please remember your child should not be in school for 48 hours after their last bout of diarrhoea or sickness. This reduces the risk of the bug spreading around the school to pupils especially any vulnerable pupils and staff.

Here is further useful material to maintain good health over the coming months and some clarity of school practice.

Thank you for your continued support

Hydration:

At Eastlands, we encourage every pupil to bring a water bottle to school with them to maintain their hydration levels.

Why?

Recent studies suggest that adequately

hydrated children may perform better in school as they are able to maintain their focus and attention throughout the entirety of the school day.

What should we drink?

As the infographic shows, the best liquid to be consumed regularly is water as it hydrates without providing extra energy (calories) or harming teeth.


As a result of this information, we insist that children arrive at school equipped with a water bottle that contains WATER ONLY.

Healthy Snacks:

During break, we have implemented a 'healthy snacking policy'. This requires pupils to bring a healthy snack to school to be consumed during break.

Why?

Too much sugar can lead to harmful fat building up inside and serious health problems, including painful tooth decay. Therefore, fruit and vegetables are always the best choice for a break time snack.



Safeguarding is Everybody's Responsibility Eastlands Primary School Lansdowne Place, Rugby CV21 3RY Tel: 01788 57328



Preparing for



Eastlands Primary School

PE Health & Well-being

Daily Mile:

uccess in a changing world.

Since last summer term, Eastlands pupils have been participating in 'The Daily Mile'; a 15 minute walk or jog at the end of every lunch.

Why?

Research has shown that participating in 'The Daily Mile' has huge health and educational benefits, including:

An increase in Fitness, stamina and energy levels.

An improvement in body composition – bone density, muscle strength, joints and cardiovascular health

Development in gross and fine motor skills, leading to better balance.

Exercise:

As a school, it is a <u>requirement</u> that we provide a curriculum that offers pupils with the opportunity to be physically active throughout the day. Each day, a pupil should be physically active for a minimum of 30 minutes a day.

But, what is actually recommended?

It is <u>recommended</u> by the Department of Health and Social Care that a child over the age of 5 should aim to be active for a minimum of 60 minutes per day (on average); whilst a child of age 5 or under should aim to be active for a minimum of 180 minutes (3 hours) a day. The images provide suggestions about raising physical activity both inside and outside of school.

What's New at Eastlands?

What an amazing start to the year we have had! Since the beginning of the year, we have rolled out a new approach to PE lesson in Real PE—a new scheme that focuses on the fundamental skills of Physical Activity as opposed to sport specific lessons! This ensures that every PE lesson is 100% inclusive and accessible to all abilities. Additionally, we welcome Mr. Batchelor to the Eastlands family; Mr. Batchelor will be supporting the pupils throughout every PE lessons and putting on some exciting new activities during lunchtimes.

Useful websites:

https://www.nhs.uk/change4life https://www.bbc.co.uk/teach/supermovers https://www.gonoodle.com/

https://thedailymile.co.uk/

Safeguarding is Everybody's Responsibility Eastlands Primary School Lansdowne Place, Rugby CV21 3RY Tel: 01788 57328





Eastlands Primary School Protective Behaviours Curriculum



🖕 mpowering &

Preparing for

Duccess in a changing world.

Eastlands Primary School

Warwickshire School Health and Wellbeing Service

Your school nurse led service is friendly, confidential and free for all school aged children (aged 5 – 19 years*) and their families. We can offer you and your child support on a wide range of health related issues such as:



Child Development e.g. bedwetting, sleep, diet Relationships Physical Health Healthy Eating & Lifestyle Choices Emotional & Mental Health Bullying Managing Medicines Childhood Immunisations Smoking Alcohol & Drugs Sexual Health

If your child has a learning need, disability or long term health condition (e.g. epilepsy, asthma, diabetes, allergies) we can help you get the right care and support.

Friendly

Helpful

Safe

*up to 25 years for young people with special educational needs or disabilities.

For more information about this service please contact the team on: 03300 245 204 or warwickshireSH&WBService@compass-uk.org





We all have the right to feel safe all the time. Eastlands Primary School Protective Behaviours Curriculum

March 2017 Public Health visit www.publichealth.hscni.net or www.gov.uk/government/organisations/Public-health-england if you would like any further advice or information, including the latest guidance. Children with rashes should Agency be considered infectious and assessed by their doctor schools and other childcare settings Juidance on infection control in Prevent the spread of infections by ensuring: routine immunisation, high standards of personal

hygiene and practice, particularly handwashing, and maintaining a clean environment. Please contact the Public Health Agency Health Protection Duty Room (Duty Room) on 0300 555 0119 or

Press and a		
Kashes and skin infections	Recommended period to be kept away from school, nursery or childminders	Comments
Athlete's foot	None	Athlete's foot is not a serious condition. Treatment is recommended
Chickenpox*	Until all vesides have crusted over	See: Vulnerable children and female staff - pregnancy
Cold sores, (Herpes simplex)	None	Avoid kissing and contact with the sores. Cold sores are generally mild and self-limiting
German measles (rubella)*	Four days from onset of rash (as per "Green Book")	Preventable by immunisation (MMR x 2 doses). See: Female staff – pregnancy
Hand, foot and mouth	None	Contact the Duty Room if a large number of children are affected Exclusion may be considered in some circumstances
Impetigo	Until lesions are crusted and healed, or 48 hours after commencing antibiotic treatment	Antibiotic treatment speeds healing and reduces the infectious period
Measles*	Four days from onset of rash	Preventable by vaccination (MMR x 2). See:Vulnerable children and female staff – pregnancy
Molluscum contagiosum	None	A self-limiting condition
Ringworm	Exclusion not usually required	Treatment is required
Roseola (infantum)	None	None
Scabies	Child can return after first treatment	Household and close contacts require treatment
Scarlet fever*	Child can return 24 hours after commencing appropriate antibiotic treatment	Antibiotic treatment recommended for the affected child if more than one child has scarlet fever contact PHA Duty Room for further advice
Stapped cheek (firth disease or parvovirus B19)	None once rash has developed	See-Vulnerable children and female staff – pregnancy
Shingles	Exclude only if rash is weeping and cannot be covered	Can cause chickenpox in those who are not immune the have not had chickenpox. It is spead by we does contact and touch if further information is required, contact the Duty Room SEE vulnerable Children and Fendle Staff – Pregnancy
Warts and verrucae	None	Vernucae should be covered in swimming pools, gymnasiums and changing rooms

We all have the right to feel safe all the time. **Eastlands Primary School Protective Behaviours Curriculum**

Good hypitiene practice Handwalling is one of the most important ways of controlling the spread of infections, especially those that cause diarrhoes and vormiting, and requiratory disease. The recommended method is the use of liquid scap, warm waters and paper tawels. Always weah hands after using the tolet, before earling or handling food, and tech handling archimed. Force all outs and disordiner with waterproof diseasings.

Coupling and streeting easily spread intections. Unliden and soluts should be encouraged to cover their mouth and nose with a tissue. Weah hands of a discouraged or discouraged.

Personal protective equipment (PPE). Dispossible non-powdered vityl or lates: free CE-marked glowes and disposable plastic aprova must be wom where there is a risk of splashing or contamination with blood/hody fluids (for example, narpy) or pad changing). Goggles should also be available for use if there is a risk of splashing to the face. Correct PFE should be used when handling clearing chemicals.

Cleaning of the environment, including toys and equipment, should be frequent, through and follow national guidance. For example, use colour-equipment, follow Control of Substances Harardous to Health (COSHI) regulations and correct decontamination of cleaning equipment. Monitor cleaning contracts and ensure dearens are appropriately trained with access to PR.

Cleaning of blood and body fluid spillages. All yillages of blood, facess, salica vornit, rasal and ge discharges should be cleared up immediately (shargy were PL) When yillages cour, clean using a pould that complex both a deferrat and a district. Use use transmitschare's instructions and ensure its deferre against brainers and vinness and antable for use on the effectual attrace. New ensure most of clearing up blood and body fluid spillages – use disposable paper toweks and discusd before a combe before. A spillage list should be available for blood spills.

Laundry should be dealt with in a separate dedicated foxiby. Solied linen should be washed separately at the hottest wash the fabric will tolerate Wear PPR when handling solied linen. Children's solied dothing should be bagged to go home, never rineed by hand

Eastlands Primary School

Clinical waste. Always segregate dometic and clinical wate, in accordance with local policy. Used nappie/pack, giver, aprova and solied dressings should be stored in correct clinical waste bags in foot-operated bins. All clinical waste must be removed by a registered waste contractor. All clinical was bags should be less than two-thirds full and stored in a dedicated, score area while awaing collection.

Sharps, eg needles, should be discarded straight into a sharps bin conforming to 85 7320 and UN 3291 standards. Sharps bins must be kept off the floor (preferably wall-mounted) and out of reach of children

Sharps injuries and bites

it sins is broken as a result of a used needle injusy or blue, recourage the wound to bleed/wash thoroughly using soap and water. Contact CP or occopedinal health or go to ALE immediately. Israue local policy is in place for staff to follow. Contact the Duty Room for advice, if unsure

unimals

formake may carry intercions, so wash hands after handling animals. Health and Safety Executive for Northern Ineland (HSENI) guidelines for protecting the health and safety of children should be followed.

Animals in school (permanent or visiting). Fraure animals' living quarters are lept clean and away from food areas. Water should be disposed of regularly, and littler bases not accessible to children. Children should be thoughly deared after used registration and are performed and the supervised atter contact with similar and the ease where visiting animality have been test should be thoughly deared after use Veteriany advice ahoudd be supported around with similar animal same and the solution of the amina as a per fixedite are not submide a percharany appectie around a strain family and have and the aminal as a per fixedites are not submide a percharany appecties. In all species carry

Preparing for

Juccess in a changing world.

We all have the right to feel safe all the time.

Eastlands Primary School Protective Behaviours Curriculum

Diarrhoea and vomiting illness	Recommended period to be kept away from school, nursery or childminders	Comments
Diarrhoea and/or vomiting	48 hours from last episode of diarrhoea or vomiting	
E.coli 0157 VTEC*	Should be excluded for 48 hours from the last episode of diarrhoea	Further exclusion is required for young children ur five and those who have difficulty in adhering to hygiene practices
Typhoid* [and peratyphoid*] (enteric fever) chantus	Further exclusion may be required for some children until they are no longer excreting	Children in these categories should be excluded un there is widence of microbiological clearance. This guidance may also apply to some contacts of cases who may require microbiological clearance
(dysentery)		Please consult the Duty Room for further advice
Cryptosporidiosis*	Exclude for 48 hours from the last episode of diarrhoea	Exclusion from swimming is advisable for two we after the diarrhoea has settled
Respiratory infections	Recommended period to be kept away from school, nursery or childminders	Comments
Flu (influenza)	Until recovered	See-Vulnerable children
Tuberculosis*	Always consult the Duty Room	Requires protonged close contact for spread
Whooping cough* (pertussis)	48 hours from commencing antibiotic treatment, or 21 days from onset of illness if no antibiotic treatment	Preventable by vaconation. After treatment, non- infectious coughing may continue for many weeks. Duty Room will organise any contact tracing neces
Unter Infections	Recommended period to be kept away from school, nursery or childminders	Comments
Conjunctivitis	None	If an outbreak/cluster occurs, consult the Duty Ro
Diphtheria *	Brdusion is essential. Always consult with the Duty Room	Family contacts must be excluded until cleared to return by the Duty Room. Preventable by vaccination. The Duty Room will organise any contact tracing necessary
Glandular tever	None	
Head lice	None	Treatment is recommended only in cases where li lice have been seen
Hepatitis A*	Exclude until seven days after onset of Jaundice (or seven days after symptom onset if no Jaundice)	The duty room will advise on any vaccination or other control measure that are needed for close contracts of a single case of hepatitisA and for suspected outbreaks.
Hepetitis B*, C, HIV/AIDS	None	Hepatitis B and C and HIV are bloodbome vhuses th are not infectious through casual contact. For cleanin body fluid spills. SEE: Good Hyglene Practice
Meningoccal meningits*/ septiceenia*	Until recovered	Some forms of meningcocccal disease are preventable vaccration (see immunisation schedule). There is no re- to exclude schings or other dose contacts of a crace. In of an outbreak, it may be necessary to provide artibiot with or without meningcocccal vaccination to dose contacts. The Duty Room will advise on any action me

Asils to farms. For more information see https://www.hxeni.gov.uk/publications/perventing.or-controlling.ill-health-animal-contact-visitor-attractions

/ulnerable children

Some medical conditions make children vulnenske to infections that would nextly be serious in most children, these include those being treated for leakernis or other cances, on high dozes of steroids and with conditions that seriously reduce immunity. Schools and numeries and childraniefers will normally have been made aware of such children medical and seriously reduce information, messites and providus BP and if reposed to these, the parent/care should be information approximation should be childrengon, messites and parvians BP and if reposed to other of these, the parent/care should be information approximation should be able to the children in have stored immunisations in the same mane. The guidance is designed to give general advice to achieve and children extings. Some vulnenskie children may need further precautions to be taken, which should be discussed with the parent or care in conjunction with their medical team and school headh.

Female staff* - pregnancy

it's prograt women devidurs a reak or is in direct contact with someone with a potentially infection reach, this should be investigated by a doctor who can contact the day room for further advice. The greatest risk to pregrat women from such infections corner from their ann child/children, rather than the workplace. • Chickenbers can affect the emenance if a women has not stready had the infection. Resourt encourse to midwite and CP at av stase of resenance.

Cicklerper can affect the pregnarcy if a woman has not dready had the infection. Report exposure to midwife and GP at any stage of pregnancy. The GP and antenatal caret will arrange a blood test to check for immunity. Singlets is caused by the same virus as chickenpar, so aryone who has not had chickenpau is potentially witneetable to the infection if they have close contact with a care of shingles.

끮

German measles (tabelia). If a pregnant woman comes into contact with german measles she should inform her GP and antendati carer immediate to ercaure investigation. The infection may affect the developing baby if the woman is not immune and is asposed in sub pregnancy.

Stapped cheek dissue (fifth dissue or paroxins 819) can occasionally affect an unbom child. If exposed enly in pregnancy (before 20 weeks) inform whoever is giving antenatal care as this must be investigated promptly.

Measles during pregnancy can result in early delivery or even loss of the baby. If a pregnant woman is exposed zhe zhould immediately inform whose is giving antenated care to ensure investigation.

All female staff bom after 1970 working with young children are advised to ensure they have had two doses of MMR vaccine.

¹The above advice also applies to pregnant students.

Immunisations

린 등

Immaistion status should shays be dredied at school entry and at the time of any vascination. Parents should be encouraged to have their child immaised and any immunisation missed or further catch-up does organised through the child's CP.

for the most up to date immunisation whice and current schedule visit www.publichealth.horninet or the school health service can advise on the new national immunisation exhedula

latest national immunisation schedule.	sation schedule.	
When to Immunise	Diseases vaccine protects against	How it is given
2 months old	Diphtherta, tatanus, pertussis (whooping cough), polio and Hib	One Injection
	Pneumococcal Infection	One Injection
	Rotavirus	Orally
	Meningococcal B infection	One Injection
3 months old	Diphtherta, tatanus, pertussis, polio and Hib	One Injection
	Rotavinus	Orally
4 months old	Diphtheria, tetanus, pertussis, polio and Hib	One Injection
	Pneumococcal Infection	One Injection
	Meningococcal B Infection	One Injection
Just after the	Measles, mumps and rubella	One Injection
	Proumococcal Infection	One Injection
	Hib and meningococcal C Infection	One Injection
	Meningococcal B infection	One Injection
Every year from 2 years old up to P7	Influenza	Nasal spray or Injection

2

eded.

at Ng of

Meringliss due to other Unitil recovered becteria	Until recovered	Hb and pneumococcal meningits are preventable by vaccination. There is no reason to exclude siblings or other close contacts of a crase. The Dury Room will give advice on any action needed
Meningitis vital"	None	Milder illness. There is no reason to exclude slbings and other dose contacts of a case. Contact tracing is not required
NRSA	None	Cood hygene, in particular handwashing and environmental cleaning, are important to minimise any danger of spread. If further information is required, contact the Dury Noom
Mumps*	Exclude child for five days after onset of swelling	Preventable by vaccination (MMR x 2 doses)
Threadworms	None	Treatment is recommended for the child and household contacts
Tonsilitis	None	There are many causes, but most cases are due to viruses and do not need an antibiotic
denotes a notifiable disease.	denotes a notifiable disease. It is a statutory notinement that doctors roort a notifiable disease to the Director of Public Health via the Dury Room.	to the Director of Public Health Via the Duty Room.
Outbreaks: if a school, nutskry o	Outbreaks: if a school, nursery or childminder suspects an outbreak of infectious disease, they should inform the Dudy Room.	s inform the Duty Room.

3 years and 4	Diphtherta, tetanus, pertussis and polio	One Injection	
	Mezske, mumps and rubella	One Injection	
GHs 12 to 13 years old	Cervical carcer caused by human populitymantus types 16 and 18 and genital warts caused by types 6 and 11	Two injections over size months	
14 to 18 years old	Tetanus, diphtherta and polio	One Injection	
	Meningococcal infection ACMY	One Injection	
his is the Immunisatio he most updated versi	is is the Immunisation Schedule as of July 2016. Children who present with certain risk factors may require addition is most updated version of the "Green Book" for the latest immunisation schedule on www.gov.uk/gov.erment/colle	ors may require addit gov.uk/gov.ernment/	. <u>5</u> 5

is it the Immunisation Schedule as of July 2016. Childhen who present with certain risk factors may require softlional immunisation. Always consult runst updated version of the "Green Book" for the latest immunisation schedule on www.gouuk/government/collections/immunisation-againstections-deesse-the-green-book#the-green-book un Oncher 2017 Arithden will service hearenise in 2-3 and 4 monthe of sea in rombination with the dishthenis Internations activities color

om October 2017 children will receive hepatitis 8 vaccine at 2, 3, and 4 months of age in combination with the diphtheria, tetarux, pertuzsix, polio of Hb vaccine. lat Immunisations. All staff should undergo a full occupational health check prior to employment: this includes ensuing they are up to date with immunisations, including two doese of MMR.

Criginal material was produced by the Health Protection Agency and this version adapted by the Public Health Agency, 12.221 Linenhall Street, Belfast, B12 885. Tet 0300 555 0114. www.publichealth.Iterri.net

Eastlands Primary School

repaining for

Juccess in a changing world.

Preparing for