



## November 2022 Bulletin 2—PE Health & Well-being

Here is some useful material to maintain good health over the coming months and some clarity of school practice.

Thank you for your continued support and we hope you find this information useful.

## **Hydration:**

At Eastlands, we encourage every pupil to bring a water bottle to school to maintain their hydration levels. We recommend students to drink water after break time and lunch times and throughout the day especially with their lunch meal.


<u>Why?</u> Recent studies suggest that adequately hydrated children may perform better in school as they are able to maintain their focus and attention throughout the entirety of the school day. It is also very beneficial to the human body as we need water regularly to survive and provide for our body. Children's bodies need to replenish the water being lost through sweat ,breathing and urination and water is the best fluid for this.

Here are some of are student hydration Heroes around the school!









## PE Health & Well-being

### **Healthy Snacks:**

During break, we have a 'healthy snacking policy'. This requires pupils in Years 3 –6 to bring a healthy snack to school to be consumed during break.

Fruit/ Veg is provided, under the Government scheme, for pupils in Reception and Years 1&2.



## Why?

Too much sugar can lead to harmful fat building up inside and serious health problems, including painful tooth decay. Therefore, fruit and vegetables are always the best choice for a break time snack. Some studies have found that children with high sugar diets are more likely to engage in conflicts, have behavioural issues at school, and perform relatively poorly on standardized tests.

<u>Concerning fact:</u> Half the sugar children are having comes from snacks and sugary drinks. On average, that's the equivalent to 2,500 sugar cubes a year! We strive at Eastland's primary school to maintain healthy habits so students keep these good habits as they pass through primary school and on to their future pushing towards a healthy lifestyle.

### **Exercise:**

As a school, it is a requirement that we provide a curriculum that offers pupils with the opportunity to be physically active throughout the day. Each day, a pupil should be physically active for a minimum of 30 minutes. We advise students partake in exercise of some kind for 30 minutes a day outside of school where possible also, whether it's a walk to the park or



going to a sports club outside of school. We also have extra sports activities at lunch time and after school such as netball, football, and so on. This is so students get a wider range of sports to enjoy in there free time and also to have fun whilst exercising.


### But, what is actually recommended?

It is recommended by the Department of Health and Social Care that a child over the age of 5 should aim to be active for a minimum of 60 minutes per day (on average); whilst a child of age 5 or under should aim to be active for a minimum of 180 minutes (3 hours) a day. The images, on the following pages, provide suggestions about raising physical activity both inside and outside of school.

## Safeguarding is Everybody's Responsibility

Eastlands Primary School Lansdowne Place, Rugby CV21 3RY Tel: 01788 57328





We all have the right to feel safe all the time.



## KS1 and KS2 PE lessons!



Real PE in Eastland's primary school - real PE is a unique, child centred approach that transforms how we teach PE to include, challenge and support your child. We are part of the movement and have joined over 2,000 real PE member schools and community of 37,000 teachers creating positive relationships with physical activity for life for every child.

Real PE helps support every child to develop the physical literacy, emotional and thinking skills to achieve in PE, Sport and life. Our unique, child centred approach transforms how we teach PE to include, challenge and support every child.

It is fully aligned to the National Curriculum, Ofsted, proposed Curriculum for Wales and Estyn requirements and focuses on the development of agility, balance and coordination, healthy competition and cooperative learning through a unique and market leading approach to teaching and learning in PE.



Real PE is taught across the whole school from nursery to year 6 and there are many different types of lessons and activities beneficial to your child's needs and requirement's for there age group. Here are a couple of real PE photos.





We all have the right to feel safe all the time.



# Warwickshire School Health and Wellbeing Service

Your school nurse led service is friendly, confidential and free for all school aged children (aged 5 – 19 years\*) and their families. We can offer you and your child support on a wide range of health related issues such as:



Child Development
e.g. bedwetting, sleep, diet
Relationships
Physical Health
Healthy Eating & Lifestyle Choices
Emotional & Mental Health
Bullying
Managing Medicines
Childhood Immunisations
Smoking
Alcohol & Drugs
Sexual Health

If your child has a learning need, disability or long term health condition (e.g. epilepsy, asthma, diabetes, allergies) we can help you get the right care and support.

## **Friendly**

Helpful

Safe

\*up to 25 years for young people with special educational needs or disabilities.

For more information about this service please contact the team on: 03300 245 204 or

warwickshireSH&WBService@compass-uk.org





We all have the right to feel safe all the time.



hygiene and practice, particularly handwashing, and maintaining a clean environment. Please contact the Public Health Agency Health Protection Duty Room (Duty Room) on 0300 555 0119 or Prevent the spread of infections by ensuring: routine immunisation, high standards of personal

schools and other childcare settings

**Suidance on infection control in** 

visit www.publichealth.hscni.net or www.gov.uk/government/organisations/Public-health-england if you would like any further advice or information, including the latest guidance. Children with rashes should be considered infectious and assessed by their doctor

Good hygiene practice Handwashing to no for most invocant ways of controlling the spread of infections, especially those that cause diarchess and vomiting, and requirable disease. The ecommended method is the use of liquid soap, warm water and paper towerk. Always wash hands after using the tolled, before estings or handing into and atter handing animize. Cover all outs and abstrations with waterproof descripes.

Coughing and sneezing easily spread infections. Children and adults should be encouraged to cover their mouth and nose with a tissue. Wash hands after using or disposing of tissues. Spitting should be discouraged.

See: Vulnerable children and female staff – pregnancy

Until all vesides have crusted over

None

None

skin infections Rashes and

Athlete's foot is not a serious condition. Treatment is

Personal protective equipment (PPE). Disposable non-powdered virel or attacker. CE-marked gloves and disposable plastic aprox must be wom where there is a risk of splashing or contamination with bloodfoody fluids (for example, nappy or pad charging). Goggles should also be available for use if there is a risk of splashing to the face. Cornect PPE should be used when handling chaning chemicals.

Cleaning of the environment, including toys and equipment, should be frequent, through and follow national guidance. For example, use colour-equipment, follow Control of Substances Hazardous to Health (COSHH) regulations and cornect decontamination of cleaning equipment. Monitor cleaning contracts and ensure cleaners are appropriately trained with access to PPE.

Cleaning of bood and body fluid spillages. All spillages of blood, faces, saliva, vornit, naral and eye dischages should be cleared up immediately (slways west PRE). When spillages court, chan using a product that complice both a delegate and a discharal. Use super manifesturers instructions and ensure it is effectual using a product that complice for use on the affected surface. Never use mosp for clearing up belong and body fluid spillages – use disposable paper towerks and dischard limital waste as described below. A spillage it should be available for blood spills.

Laundry should be dealt with in a separate dedicated foxiby. Solide linen should be washed separately at the hottest wash the fabric will tolerate. Wear PPE when handling solled linen. Children's soiled clothing should be bagged to go home, never rinsed by hand.

See: Vulnerable children and female staff – pregnancy

A self-limiting condition Treatment is required

Preventable by vaccination (MMR x 2).

Antibiotic treatment speeds healing and reduces the

Until lesions are crusted and healed, or 48 hours after commencing antibiotic treatment

Contact the Duty Room if a large number of children are affected. Exclusion may be considered in some

Preventable by immunisation (MMR x 2 doses). See: Female staff – pregnancy

Four days from onset of rash (as per "Green Book")

Cold sores are generally mild and self-limiting Avoid kissing and contact with the sores.

Clinical Waste. Always segregate domestic and clinical waste, in accordance with local policy, Used nappies/pads, gloves, aprone and soiled dressings should be stored in correct clinical waste large in foot-operated birs. All clinical wast enust be removed by a registered waste contractor. All clinical wast bags should be less than two-thirds full and stored in a dedicated, secure

Sharps, eg needles, should be discarded straight into a sharps bin conforming to 85 7320 and UN 3291 standards. Sharps bins must be kept off the floor (preferably wall-mounted) and out of reach of children

See: Vulnerable children and female staff – pregnancy

None once rash has developed

child if more than one child has scarlet fever contact PHA Duty Room for further advice Antibiotic treatment recommended for the affected Household and close contacts require treatment

Child can return 24 hours after commencing

Child can return after first treatment

Exclusion not usually required

None

Can cause chickenpox in those who are not immune

Exclude only if rash is weeping and cannot be covered

i.e. have not had chickenpox. It is spread by very dose contact the Duty Room. SEE: Vulnerable Children and Female Staff – Pregnancy

Vertucae should be covered in swimming pools,

gymnasiums and changing rooms

if skin is kretien as a result of a used needle injuny or bits, encourage the wound to bleedwach thoroughly using soap and water. Contact GP or compational health or go to A&E immediately, fensue local policy is in place for staff to follow. Contact the Duty Room for advice, if ursure.

loimás may cany intections, so wach hands after handing animás. Health and Safety Executive for Northern Ireland (HSENI) guidelines for protecting the health and safety of children should be followed.

Anlinais in school (permanent or visiting), frauer animals' bring quarters are kept clean and away from food areas. Waste ahould be disposed of regularly, and litter boxes not accessable to children. Children should not play with animals unsupervised. Hand-hygiene should be supervised after contact with animals and the area where visiting animals have been tept should be thoroughly dearned after use. Veterinary advice should be sought on animal welfare and animal health issues and the suitability of the animal as a pet. Regolles are not suitable as peets in schools and runseries, as all species carry

We all have the right to feel safe all the time.

One injection One injection

Orally

Diphtheria, totanus, portussis, polio and Hib

Maningococcal B Infaction

Diphtheria, totanus, portussis, polio and Hib

Hib and moningococcal C Infaction

Meningococcal B Infection

Measles, mumps and rubella

umococcal Infection

laningococcal B Infaction

mococcal Infection



# **Eastlands Primary School**

MS. For more information see https://www.hoeni.gov.uk/publications/preventing.or-controlling.ill-health-animal-contact-visitor-attractions

# nerable childre

e medical conditions make children volverable to infections that would nately be serious in most children, these include those being treated for errains or other careces, on high dozes of sternids and with conditions that seriously reduce immunity. Schools and nurseries and children-less will also have been moste aware of such children. These children are particularly vulnerable to children proceed and parrowins 1919 and, if exposed there of these, the parent/choiner broughd be informed promptly and further medical loss coaght. If may be abhabble for these children to have tional immunications for example preumococcal and influence. This guidance is designed to give general advice to schools and children settings evulnerable children may need further precadions to be taken, which should be discussed with the parent or carer in conjunction with their real team and school health.

# ale staff\* - pregnand

FUTIONE STATE - IN FIGURATION Is pregrat woman develops a rach or is in direct contact with someone with a potentially infectious rach, this should be investigated by a doctor who can contact he day room for futther salvice. The greatest risk to pregrat women from such infections comes from their own childchildcen, rather than the workplace.

idenpor can affect the pregnacy if a woman has not already had the infection. Report exposure to midwife and CP at any stage of pregnancy, e CP and antenated care will armange a blood test to check for immunity. Shrigles is caused by the same virus as chickenpore, so anyone who has t had chickenpor is potentially vulnerable to the infection if they have close contact with a case of shingles.

ieman meades (ndella). If a pregnant women comes into contact with german meades she abould inform her CP and antenatal carer immediate o ensure investigation. The infection may affect the developing baby if the woman is not immune and is exposed in easily pregnancy.

Apped cheel dessae (lith disease or parovins 819) can occasionally affect an unioum child if exposed early in pregnancy (before 20 weeks), informwhoerer is giving anternatal care as this must be investigated promptly.

Measles during programy; can result in early delivery or even loss of the baby. If a pregnant woman is exposed she should immediately inform theorems is mission extension one to secure immediately.

female staff born after 1970 working with young children are advised to ensure they have had awo doses of MMR vaccine.

shows advice also annies to necessari stude

## estione

nisation status should always be checked at school entry and at the time of any vaccination. Barents should be encounged to have their child rised and any immunisation missed or further cutch-up dozes organised through the child's CP.

for the most up-to-date immunisation advice and current schedule visit www.publichealth.froni.net or the school health service can advice on the

Diphtheria, totanus, pertussis (whooping cough), polio and Hib

aumococcal Infection

Diseases vaccine protects against

Diarrhoea and	Recommended period to be kept away from school, nursery or childminders	Comments	Visits to fam
Diarrhoea and/or vomiting	48 hours from last episode of diarrhoea or vomiting		Vulnerabl Some medical
E.coii 0157 VTEC*	Should be excluded for 48 hours from the last episode of diarrhoea	Further exclusion is required for young children under five and those who have difficulty in adhering to hygiene practices	normally have to either of th additional imp
Typhoid" [and paratyphoid"] (enteric rever)	Further exclusion may be required for some children until they are no longer excreting	Children in these categories should be excluded until there is evidence of microbiological clearance. This guidence may also apply to some contacts of cases who may require microbiological clearance.	medical team Female Si
Shigella* (dysentery)		Please consult the Duty Room for further advice	the duty room  Olidenpax
Cryptosporidiosis*	Exclude for 48 hours from the last episode of diarrhoea	Exclusion from swimming is advisable for two weeks after the diarrhoea has settled	The GP and not had chii
Respiratory infections	Recommended period to be kept away from school, nursery or childminders	Comments	to ensure in Slapped che informwho
Flu (influenza)	Until recovered	See-Vulnerable children	Measles dur
Tuberculosis*	Always consult the Duty Room	Requires prolonged close contact for spread	whoever is g
Whooping cough* (pertussis)	48 hours from commencing artibiotic treatment, or 21 days from orset of illness if no artibiotic treatment	Peventable by vaccination. After treatment, non- infectious coughing may continue for many weeks. The Duty Room will organise any contact tracing necessary	*The above ad Immunisa
			Immunisation
Other infections	Recommended period to be kept away from school, nursery or childminders	Comments	immunised an
Conjunctivitis	None	if an outbreak/cluster occurs, consult the Duty Room	For the most of latest national
Diphtheria *	Exclusion is essential. Aways consult with the Duty Room	Pamily contacts must be excluded until cleared to return by the Duty Room.  Preventable by vaccination. The Duty Room will organise any contact tracing necessary.	When to imn 2 months old
Glandular fever	None		
Head lice	None	Treatment is recommended only in cases where live lice have been seen	3 months old
Hepatitis A*	Exclude until seven days after onset of Jaundice (or seven days after symptom onset if no Jaundice)	The duty room will advise on any vaccination or other control measure that are needed for close contacts of a single case of hepatitis A and for suspected outbreals.	4 months old
Hepatitis B*, C, HTV/AIDS	None	Hepatitis B and C and HIV are bloodhome viruses that are not infectious through casual contact. For cleaning of body fluid spills. SEE Good Hygene Practice	Just after the
Meningboocal meningbb*/ septicaemia*	Until recovered	Some forms of meningococcal desease are preventable by vaccination (see immunication schedule). There is no neason to enclude splings or other dose contacts of a case. In case of an outbreak, it may be necessary to provide artibiotics with or without meningococcal vaccination to dose contacts. The Doty Room will advise on any action needed.	Every year fr years old up

We all have the right to feel safe all the time.



3 years and 4	Diphtherta, totanus, pertussis and polio	One Injection
III OIL	Mezales, mumps and rubella	One Injection
GHs 12 to 13 yvars old	Genvical cancer caused by human papillomavitus types 15 and 18 and genital warts caused by types 6 and 11	Two injections over six months
14 to 18 years old	Tetanus, diphthenta and polio	One Injection
	Maningococcal Infaction ACVY	One Injection

Milder illness. There is no reason to exclude siblings and other close contacts of a case. Contact tracing is

Aeningitis viral\*

give advice on any action needed

ervironmental cleaning, are important to minimise

any danger of spread. If further information is required, contact the Duty Room

Preventable by vaccination (MMR x 2 doses)

Exclude child for five days after onset of

Good hygiene, in particular handwashing and

#SA

Hib and pneumococcal meningitis are preventable b vaccination. There is no reason to exclude siblings or other close contacts of a case. The Duty Room will his is the Immunisation Schedule as of July 2016. Children who present with certain risk factors may require additional immunisations. Always consult the most updated version of the "Green Book" for the latest immunisation schedule on www.gocub/gaverment/collections/immunisation-against-infectious-diseases the green-booksfate-green-book

rom October 2017 children will receive hepatitis 8 vaccine at 2,3, and 4 months of age in combination with the diphtheria, tetanus, pertussis, polio and Hib vaccine. Staff immunisations, All staff should undergo a full occupational health check prior to employment; this includes ensuring they are up to date with immunisations, including two doses of MMR. Driginal material was produced by the Health Protection Agency and this version adapted by the Public Health Agency,

There are many causes, but most cases are due to

viruses and do not need an antibiotic

Treatment is recommended for the child and

household contacts

12-22 Linenhall Street, Belfast, 872 88S. Tek 0300 555 0114.

nformation produced with the assistance of the Royal College of Paediatrics and Child Health and Public Health England ww.publichealth.hsmi.net

denotes a notifiable classure. It is a statutory requirement that doctors report a notifiable classor to the Director of Public Health via the Duty Room Outbreaks: if a school, rursary or childminder suspects an outbreak of intectious disease, they should inform the Duly Room

We all have the right to feel safe all the time.